

# CONDITION FACTOR OF THREE SPECIES FROM THE BULGARIAN BLACK SEA COAST

Maria YANKOVA<sup>1</sup>

<sup>1</sup>Institute of Oceanology, BAS, Varna, Bulgaria, Asparuhovo quarter 40, First of May str. Corresponding author Maria Yankova, e-mail: maria\_y@abv.bg

**Abstract:** The condition factors of *Trachurus mediterraneus ponticus*, *Engraulis encrasicolus* and *Sarda sarda* were studied. Mean condition factor for *T. mediterraneus ponticus* was  $0.77 \pm 0.28$ , while it was  $0.63 \pm 0.17$  and  $2.47 \pm 1.43$  for *E. encrasicolus* and *S. sarda*, respectively. There was a significant difference ( $p < 0.05$ ) in the mean condition factor of male and female *T. mediterraneus ponticus*, while in *S. sarda* and *E. encrasicolus* there was not a significant difference in the condition factors of males and females. It was observed that the autumn season condition factors of *T. mediterraneus ponticus* and *E. encrasicolus* were higher than summer season values.

**Key Words:** condition factor, horse mackerel, anchovy, Atlantic bonito, Black Sea

## I. INTRODUCTION

The mid-water pelagic-trawl commercial target species in the Bulgarian Black Sea study area are small pelagic fishes, mainly anchovy and horse mackerel. The values of the local catches are relatively high, as a consequence of the high demand for these resources for direct consumption (Yankova 2013b). The small pelagic-species production is characterized by high interannual fluctuations that appear to be unrelated to the level of fishing effort. On the contrary, there is strong evidence that yearly production is connected to the success of reproduction, which in turn can be related to the direct or indirect effect of favorable environmental conditions (i.e. the availability of food, as determined by the timing of zooplankton (Yankova *et al.* 2014). The Atlantic bonito, *Sarda sarda* (Bloch, 1793), is one of the most abundant small tuna-like species which exists with a wide ranging distribution spanning tropical and temperate areas of the Atlantic ocean, including the Mediterranean and Black Sea (Yoshida, 1980; Sabatés and Recasens, 2001).

In fisheries science, the condition factor is used in order to compare the “condition”, “fatness” or wellbeing of fish and it is based on the hypothesis that heavier fish of a particular length are in a better physiological condition (Bagenal, 1978). Condition factor is also a useful index for the monitoring of feeding intensity, age, and growth rates in fish (Oni *et al.*, 1983). It is strongly influenced by both biotic and abiotic environmental conditions and can be used as an index to assess the status of the aquatic ecosystem in which fish live. Condition factors of horse mackerel was investigated and reported by Yankova (2009a, b), Yankova and Raykov (2009), Yankova (2010), Yankova (2013a). Despite their importance, little is known about the biology and ecology of *S. sarda* and *E. encrasicolus* in the Bulgarian Black Sea waters. For instance, only a few studies have provided some biological information of the anchovy in the Bulgarian Black Sea (Mikhailov 2005). However, the condition factors of both species have not yet been analyzed. These report focused on

the determination of changes in condition factor with season, fish length, sex and reproductive status of fish.

## II. MATERIAL AND METHODS

Fish samples were collected between April 2012 and December 2012 from the commercial catches landed in Bulgarian Black Sea coast. A total of 3080 fish were collected throughout the study period. Fork length (FL) of each individual fish was measured to the nearest cm by using a measurement board. The length frequency data were pooled into groups of 0.5 cm length intervals. The sex was assigned macroscopically. Fulton's condition factor was calculated from the expression (Bagenal, 1978):  $K = 100 W/L^3$ . Where W is the whole body weight in grams and L the standard length in millimetres. All results obtained in these analyses of *T. mediterraneus*, *E. encrasicolus* and *S. sarda* are carried out for male, female and total samples.

## III. RESULTS

The mean condition factors for the *Trachurus mediterraneus ponticus*, *Engraulis encrasicolus* and *Sarda sarda* in the Bulgarian Black Sea coast are shown in Table 1. The results indicated that there was a significant difference between the condition factors of male and female *T. mediterraneus* ( $p < 0.05$ ). There were not significant differences between the condition factors of male and female fish in other two species.

Table 1. Condition Factor of *T. mediterraneus ponticus*, *E. encrasicolus* and *S. sarda* in the Bulgarian Black Sea coast.

Species	Summer	Number	Autumn	Number	T-test	P
	season		season			
<i>T. mediterraneus ponticus</i>	0.74±0.32	340	0.99±1.12	270		0.05
<i>E. encrasicolus</i>	0.68±0.29	420	0.85±0.92	290		0.05
<i>S. sarda</i>	-	-	2.53±1.49	340		0.05

The condition factors were also checked during summer and autumn seasons (Table 2). The condition factor for *T.*

### Publication History

Manuscript Received : 11 March 2015  
 Manuscript Accepted : 26 March 2015  
 Revision Received : 25 April 2015  
 Manuscript Published : 30 April 2015


*mediterraneus ponticus* in autumn season ( $0.99 \pm 1.12$ ) was significantly higher than that ( $0.74 \pm 0.32$ ) in summer season ( $p < 0.05$ ). The same trend was observed in *E. encrasicolus* while the autumn condition factor ( $0.85 \pm 0.92$ ) was significantly higher than the mean value of summer season ( $0.68 \pm 0.29$ ) ( $p < 0.05$ ).

Table 2. Seasonal variation and t-test analysis of the condition factor of both sexes of *T. mediterraneus ponticus*, *E. encrasicolus* and *S. sarda* in the Bulgarian Black Sea coast.


Species	Summer season	Number
<i>T. mediterraneus ponticus</i>	$0.74 \pm 0.32$	340
<i>E. encrasicolus</i>	$0.68 \pm 0.29$	420
<i>S. sarda</i>	-	-

Figure 2 shows the variations in condition factors with standard lengths. In *S. sarda* (Fig.2A), three phases are easily distinguished. In phase I, there was a significant increase from  $2.94 \pm 0.15$  (27 cm) to  $3.36 \pm 0.50$  at a length of 29 cm ( $p < 0.05$ ). In phase II, there was a decline to  $1.73 \pm 0.30$  at a standard length of 33 mm. In the third phase the condition factor increased from  $1.73 \pm 0.30$  at a length of 33 cm to  $2.14 \pm 0.3$  at a size of 34 cm.


In *T. mediterraneus ponticus* (Figure 2B), the size increase ranged from  $0.84 \pm 0.05$  (15 cm) to  $1.35 \pm 0.05$  (15.5 mm) ( $p < 0.01$ ). Then, it significantly slumped from the previous level ( $1.35 \pm 0.05$ ) to  $0.81 \pm 0.41$  at a length of 16 cm ( $p < 0.05$ ). Condition factor values of *E. encrasicolus* (Figure 2C), registered a significant decrease ( $p < 0.05$ ) between the size range of 13 cm and 13.5 cm from  $0.53 \pm 0.20$  to  $0.31 \pm 0.11$ , respectively. Thereafter, it significantly increased ( $p < 0.05$ ) to a mean value of  $0.58 \pm 0.22$ .



A.



B.



C.

Figure 2. Variation of condition factor with fish size.

#### IV. DISCUSSION

However, the condition factor values of *T. mediterraneus ponticus* from the current study are relatively higher than those reported for the other study (Yankova, 2009; Yankova and Raykov, 2009; Yankova, 2010). Similarly, the condition factor for *E. encrasicolus* in our study is higher than that found for the same species by Giraldez and Abad, 1995 for western Mediterranean. Similar results have been observed by Aka *et al.*, 2004 for the Black Sea. Environmental factors have been taken into consideration in order to account for spatial and temporal differences in condition factor of fish (Bagenal, 1978; Nagel, 1986; Ekrem, 2004). P

The results of the current study also presented that there were no significant variations in condition factors with respect to sex, on the contrary to *E. encrasicolus*, but the study was in agreement with the values reported for anchovy (Sinovčić, 2000; Millan, 1999; Özdamar, 1991). It was found that autumn season condition factor of *T. mediterraneus ponticus* and *E. encrasicolus* were higher than summer season values. Seasonal variation in the condition factor of fish has been reported for *T. mediterraneus ponticus*. However, the results of this study do not conform to those published for *E. encrasicolus* (Mikhailov, 2005) in which no seasonal changes were observed in condition factor. Despite this difference in observation, Oni *et al.*, (1983) noted that condition factor is not constant for a species or population over a time interval and might be influenced by both biotic and abiotic factors such as feeding regime and state of gonadal development (Saliu, 2001). Also, Alegria-Hernandez (1994) reported that changes in the condition of the fish are related to the development and maturation of the gonads when, due to the significant energy expenditure for these processes, the condition of the fish declines.

This study affirmed that there is a variation in condition factor by size (length) classes in three species studied. The general trend is that in *S. sarda* relatively lower condition factors were recorded for relatively large sizes, while for *T. mediterraneus ponticus* and *E. encrasicolus* relatively higher condition factors were recorded for rather smaller fish.

#### V. CONCLUSIONS

This study presented the basic information on the condition factors for *T. mediterraneus ponticus*, *E. encrasicolus* and *S. sarda* species from the Bulgarian Black Sea coast, which would be useful for fishery managers as well as the sustainable management of its stocks in the region. Moreover, there are no conditions factors currently in the Fish Base for these species and therefore, our results may contribute to this invaluable database.

#### REFERENCES

- [1] Aka, Z., T Hatice, Turan C. 2004. A study on the growth of the anchovy *Engraulis encrasicolus* Linnaeus (1758) In Turkish Seas. Pakistan Journal of Biological Sciences 7(7): 1121-1126
- [2] Alegria-Hernandez, V. 1994. Reproductive cycle and changes in conditions of the horse mackerel (*Trachurus trachurus* L.) from the Adriatic Sea. Acta Adriat. 35, 59-67.
- [3] Bagenal, T.B. 1978. Aspects of fish fecundity. In: S.D. Gerking (Ed) Ecology of Freshwater fish Production. Blackwell Scientific Publications, Oxford: 75-101.

- [4] Braga, F.M.S. 1986. Estudo entre o fator de condição e relação peso/comprimento para alguns peixes marinhos. Rev. Brasil. Biol., 46(2): 339-346
- [5] Ekanem, S.B. 2004. The biology and culture of the silver catfish (*Chrysichthys nigrodigitatus*). Journal of Sustainable Tropical Agricultural Research, 10: 1-7.
- [6] Giraldez, A., Abad, R. 1995. Aspects of the reproductive biology of the western Mediterranean anchovy from the coasts of Malaga (Alboran Sea). Sci. Mar., 59:15-23 pp.
- [7] Millan, M., 1999. Reproductive characteristics and condition status of anchovy *Engraulis encrasicolus* L. from The Bay of Cadiz (SW Spain), Fisheries Res., 41: 73-86.
- [8] Mikhailov, K. 2005. Some aspects of the reproductive biology of the anchovy, *Engraulis encrasicolus* (L.) off the Bulgarian Black Sea coast. Reported at the Symposium "Protection and Sustain. Manag. Black Sea, Imperative 3rd Millenium, 19-21 Oct, 2005, Constanta, Romania. Cercetari Marine.
- [9] Oni, S.K., Olayemi, J.Y., Adegboye, J.D. 1983. Comparative physiology of three ecologically distinct fresh water fishes, *Alestes nurse* Ruppell, *Synodontis schall* Bloch and *S.schneider* and *Tilapia zilli* Gervais. J. Fish Biol., 22: 105-109.
- [10] Özdamar, E. 1991. A research on estimation of some parameters of anchovy (anchovy *Engraulis encrasicolus* L. 1758) stock in the Black Sea as viewpoint of population dynamics, PhD. Thesis, Ondokuz Mayıs University, Samsun, pp:85.
- [11] Saliu, J.K. 2001. Observation on the condition factor of *Brycinus nurse* (Pisces: Cypriniformes, Characidae) from Asa Reservoir, Ilorin, Nigeria. Tropical Freshwater Biology, 10: 9–17.
- [12] Sinovčić, G. 2000. Anchovy, *Engraulis encrasicolus* (Linnaeus, 1758): biology, population dynamics and fisheries case study, Acta Adriat, 41: 3-53.
- [13] Sabatés, A., Recasens, L. 2001. Seasonal distribution and spawning of small tunas (*Auxis rochei* and *Sarda sarda*) in the north-western Mediterranean. Sci. Mar., 65(2): 95–100.
- [14] Yankova, M., Raykov, V. 2009a. Resent investigation on population structure of Horse mackerel (*Trachurus mediterraneus ponticus* Aleev., 1956) in the Bulgarian Black Sea coast. Proceedings of the Institute of Fishing Resources Varna, Volume 27, 39 - 46.
- [15] Yankova, M. 2009b. Condition factor, sex ratio and length-weight relationship, of Horse mackerel (*Trachurus mediterraneus*) from the Bulgarian Black Sea coast. Proceedings of the Union of Scientists - Varna, Series "Technical Sciences" 2'2008/1'2009, 70 - 72.
- [16] Yankova, M. 2010. Some biological aspects of the horse mackerel catch of the Bulgarian Black Sea Coast. Cercetari marine - Recherches marines, 39, NIMRD, 239 - 249.
- [17] Yankova, M. 2013a. Population Dynamics of Horse Mackerel (*Trachurus mediterraneus ponticus*) in the Bulgarian Black Sea Coast. ISRN Zoology Volume 2013, Article ID 127287, 6 pages, <http://dx.doi.org/10.1155/2013/127287>.
- [18] Yankova, M. 2013b. An overview on the biology of horse mackerel, *Trachurus mediterraneus*, off the Bulgarian Black Sea coast. In: Animal Diversity, Natural History and Conservation, (Eds.) V.K.Gupta and Anil K. Verma. Daya Publishing House, New Delhi, Vol.2, p.165-179.
- [19] Yankova, M., Stefanova, K., Doncheva, V. 2014. Influence of the marine environmental variability on the population parameters of horse mackerel (*Trachurus mediterraneus ponticus* Aleev, 1956) in front of the Bulgarian Black Sea coast. Proceedings of the Union of Scientists – Varna, Series "Marine Science" ISSN 1314-3379, 31-31.
- [20] Yoshida, H.O. 1980. Synopsis of biological data on bonitos of the genus *Sarda*. NOAA Tech. Rep. NMFS Circ. 432. FAO Fish. Synop., 118: 1-50.